

# Inhibition of Soluble Epoxide Hydrolase Limits Niacin-induced Vasodilation in Mice

Ahmet B. Inceoglu, PhD,\* Heather L. Clifton, BS,† Jun Yang, PhD,\* Christine Hegedus, PhD,\*  
Bruce D. Hammock, PhD,\* and Saul Schaefer, MD†

**Background:** The use of niacin in the treatment of dyslipidemias is limited by the common side effect of cutaneous vasodilation, commonly termed flushing. Flushing is thought to be due to release of the vasodilatory prostanoids prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) and prostaglandin E<sub>2</sub> from arachidonic acid metabolism through the cyclooxygenase pathway. Arachidonic acid is also metabolized by the cytochrome P450 system, which is regulated, in part, by the enzyme soluble epoxide hydrolase (sEH).

**Methods:** These experiments used an established murine model in which ear tissue perfusion was measured by laser Doppler to test the hypothesis that inhibition of sEH would limit niacin-induced flushing.

**Results:** Niacin-induced flushing was reduced from 506 ± 126% to 213 ± 39% in sEH knockout animals. Pharmacologic treatment with 3 structurally distinct sEH inhibitors similarly reduced flushing in a dose-dependent manner, with maximal reduction to 143% ± 15% of baseline flow using a concentration of 1 mg/kg TPAU (1-trifluoromethoxyphenyl-3-(1-acetylpiperidin-4-yl) urea). Systemically administered PGD<sub>2</sub> caused ear vasodilation, which was not changed by either pharmacologic sEH inhibition or sEH gene deletion.

**Conclusions:** Inhibition of sEH markedly reduces niacin-induced flushing in this model without an apparent effect on the response to PGD<sub>2</sub>. sEH inhibition may be a new therapeutic approach to limit flushing in humans.

**Key Words:** nicotinic acid, prostaglandins, arachidonic acid, flushing, dyslipidemia, epoxide hydrolase

(*J Cardiovasc Pharmacol*<sup>TM</sup> 2012;60:70–75)

Received for publication July 18, 2011; accepted March 29, 2012.  
From the Departments of \*Entomology; and †Internal Medicine, University of California, Davis, CA.

Supported by a UC Davis Collaborative Research Grant (to S.S.), a National Institute of Environmental Health Sciences (NIEHS) Grant R01 ES002710 (to B.D.H.), and an NIEHS Superfund Basic Research Program P42 ES004699.

The authors report no conflicts of interest.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site ([www.jcvp.org](http://www.jcvp.org)).

Reprints: Saul Schaefer, MD, Department of Internal Medicine, University of California, One Shields Avenue TB 172, Davis, CA (e-mail: [sschaefer@ucdavis.edu](mailto:sschaefer@ucdavis.edu)).

Copyright © 2012 by Lippincott Williams & Wilkins

## INTRODUCTION

Nicotinic acid, or niacin, is a water-soluble B vitamin that, in high doses, is effective in treating dyslipidemias<sup>1,2</sup> and reduces both cardiovascular mortality and cardiac events.<sup>3</sup> However, the use of niacin is limited by a major side effect, cutaneous vasodilation, which is commonly described as flushing involving the head and upper torso.<sup>4</sup> Flushing occurs in up to 90% of patients, results in discontinuation in approximately 30% of patients,<sup>5</sup> and is only marginally suppressed by interventions such as intake with food and pretreatment with cyclooxygenase (COX) inhibitors such as aspirin.<sup>6</sup> Flushing is not a cosmetic side effect but often an intense, although transient, response and a major factor for patient noncompliance that greatly reduces the potential utility of niacin. Indeed, the approval in the European Union of the D prostanoid receptor 1 (DP1) inhibitor laropiprant to limit flushing highlights the serious nature of flushing.<sup>7</sup>

A prominent mechanism of niacin-induced flushing has been described using a mouse model of niacin-induced flushing<sup>8</sup> and in cell lines and tissue samples.<sup>9</sup> Briefly, flushing is, in part, a result of niacin activating the GPR109A receptor and, through a pathway involving arachidonic acid (ARA), causing the release of the vasodilatory prostanoids prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) acting on subepidermal blood vessels. The role of PGD<sub>2</sub> as an important vasodilatory compound is supported by the efficacy of specific antagonists to the DP1 receptor to partially reduce vasodilation in mice and symptomatic flushing in man.<sup>8,10</sup> However, because these inhibitory effects are incomplete, it is likely that this proposed mechanism is an oversimplification of the flushing process and other mechanisms of niacin-induced flushing have to be elucidated.

In addition to metabolism by the COX enzymes to prostanoids, ARA is also metabolized by 2 other enzymatic routes, the lipoxygenase and cytochrome P450 (cyp450) enzymes to leukotrienes and oxygenated fatty acids, respectively. One group of products of the cyp450 pathway is epoxyoxygenated fatty acids, including epoxyeicosatrienoic acids (EETs),<sup>11</sup> which are principally regulated by rapid degradation to dihydroxyeicosatrienoic acids (DHETs) by the enzyme soluble epoxide hydrolase (sEH).<sup>12</sup> Inhibitors of sEH not only stabilize the levels of EETs but also modulate the metabolism of ARA in a manner that is not governed by rules of mass action, principally by shifting ARA flow from the COX branch toward the other 2 branches.<sup>13</sup>

Given the ability of soluble epoxide hydrolase inhibitors (sEHI) to alter AA metabolism, we hypothesized that

inhibition of sEH would also reduce niacin-induced flushing. To test this hypothesis, we used an established murine model of niacin-induced flushing, quantifying blood flow in the ear by laser Doppler flowmetry,<sup>8</sup> with niacin and sEH inhibition. These data demonstrate that genetic or pharmacologic inhibition of sEH by structurally different sEH inhibitors is highly effective in attenuating niacin-induced flushing; sEH inhibition, however, does not reduce the vasodilatory response to PGD<sub>2</sub>. Our findings strongly argue for the presence of several mechanisms that mediate flushing in mice and possibly in humans<sup>14</sup> and the need for further investigation of these mechanisms if the utility of niacin is to be expanded to a larger population of cardiovascular patients.

## MATERIALS AND METHODS

All studies were approved by the UC Davis Institutional Animal Care and Use Committee. Animals were housed under standard conditions with ad libitum food and water and a 12:12 light:dark cycle at the UC Davis facilities.

### Mouse Model

C57BL/6 male mice were obtained from Charles River Laboratories (Wilmington, MA), whereas the sEH-targeted knockout (KO) mice were used from a UC Davis colony.<sup>15</sup> For the experiments, mice were anesthetized using Nembutal (50 mg/kg) given by intraperitoneal injection. Niacin was administered subcutaneously (SC) at a concentration of 30 mg/kg in physiologic saline (equivalent to a human dose of approximately 2 g). The sEH inhibitors and other compounds (eg, aspirin, celecoxib) were administered SC over a range of relevant doses 45 minutes before niacin. PGD<sub>2</sub> (1 mg/kg) was administered SC to wild-type mice with or without TPAU 1 mg/kg and sEH  $-/-$  mice.

### Laser Doppler Ear Blood Flow

The change in ear flow was measured using a laser Doppler flow meter (BLF 21; Transonic Systems, Inc., Ithaca, NY). The laser Doppler probe was fitted with a sleeve of 2 mm length plastic tubing and attached to a micromanipulator to standardize the depth of the tissue being measured. The flow probe was placed against the ventral aspect of the right ear of the anesthetized mouse as described by Cheng et al.<sup>8</sup> Blood flow was measured at 30-second intervals before and during exposure to compounds. Baseline blood flow was established by the average of measurements over 3–5 minutes before injection of drug or vehicle. Data were analyzed as percentage of baseline blood flow in tissue perfusion units.

### Inhibitors of sEH

Structurally different sEH inhibitors were used to test the hypothesis that inhibition of flushing was a class effect of these compounds. Experiments were conducted using the sEH inhibitors TPAU (1-trifluoromethoxyphenyl-3-(1-acetylpiperidin-4-yl) urea, 0.01–1 mg/kg), t-AUCB (4 mg/kg), and sorafenib (4 mg/kg). For comparison, identical experiments were performed using aspirin (4 mg/kg) and celecoxib (4 mg/kg) to inhibit COX-1 and COX-2 pathways. The sEHs were synthesized in house,<sup>16</sup> aspirin was from Fisher

Scientific (Pittsburgh, PA), and celecoxib was from Tocris (Ellisville, MO).

### Liquid Chromatography–tandem Mass Spectrometry

A liquid chromatography–tandem mass spectrometry–based method as described earlier with modifications was used to quantify EETs and DHETs as markers of the cyp450 system and efficacy of sEH inhibition.<sup>17,18</sup> Plasma samples were obtained from cardiac puncture at 5 minutes after administration of niacin or niacin + TPAU and control. Blood was placed in a serum separator tube (BD Microtainer, Franklin Lakes, NJ), allowed to coagulate for 30 to 45 minutes, centrifuged at 10,000 revolutions per minute for 15 minutes, and then 250  $\mu$ L of plasma was extracted and stored at  $-80^{\circ}\text{C}$ . Immediately after cardiac puncture, the mouse was perfused with saline to remove any remaining blood and ears were cut, weighed, and also stored at  $-80^{\circ}\text{C}$ .

Samples were spiked with a suite of deuterium-labeled odd chain length analogs (surrogates) and then solvent was extracted and partially purified by passing through a solid-phase extraction column using Oasis HLB cartridges. The loaded column was washed with 2 mL 2.5 mM H<sub>2</sub>PO<sub>4</sub> + 10% methanol and dried under vacuum. Analytes were eluted with 2 mL of ethyl acetate. The collected extracts were evaporated to dryness under nitrogen, redissolved in a final volume of 100  $\mu$ L with methanol containing deuterated isomers of internal standards including prostaglandins, and stored at  $-80^{\circ}\text{C}$  until analysis. A 10  $\mu$ L aliquot of each sample was separated by reversed phase high-performance liquid chromatography on a 150  $\times$  2-mm internal diameter 5- $\mu$ m particle size C18 column using a gradient of water - 0.1% acetic acid and 89:11 acetonitrile to methanol (vol/vol) - 0.1% acetic acid. The separated analytes were quantified using negative mode electrospray ionization and tandem mass spectrometry in multi-reaction monitoring mode (SRM; Waters Alliance 2795 LC system and Quattro Ultima tandem-quadrupole mass spectrometer, Micromass). The system was calibrated with a minimum of 5 calibration solutions containing analytical targets at concentrations ranging from 1 to 1000 nM. The analysis of each sample was repeated 3 times. Calibration check solutions were analyzed at a minimum frequency of 10 hours to ensure stability of the analytical calibration throughout a given analysis.

### Statistical Analyses

Statistical analysis of the results was done using SigmaPlot (SigmaStat Software, Inc, Chicago, IL). Data are expressed as mean  $\pm$  standard error. Differences in the flushing response were first analyzed for normality of residuals and homogeneity of variance at an alpha level of 0.05. When these tests were not passed, the data were then analyzed by a 1-way analysis of variance (ANOVA) based on ranks (Kruskal–Wallis test) followed by a comparison of group means using Dunn method. Differences in EETs and DHETs levels were also first analyzed for normality of residuals and homogeneity of variance at an alpha level of 0.05. When these tests were passed, the data were analyzed by 1-way


ANOVA followed by Dunnett test comparing group means with the vehicle control. Significance level was set at 0.05 probability level.

**RESULTS**

**Niacin-induced Flushing Is Decreased by Genetic Deletion or Pharmacologic Inhibition of sEH**


As previously described,<sup>8</sup> niacin administration led to a significant increase in ear perfusion that peaked approximately 4 minutes after injection and returned to near control levels by 15 minutes after injection (Fig. 1). This increase in perfusion was significantly limited in sEH KO mice (ANOVA,  $P < 0.001$ ), as reflected by a decrease in maximal blood flow from  $506 \pm 126\%$  to  $213 \pm 39\%$  of baseline blood flow. Pretreatment with the sEH inhibitor TPAU (1 mg/kg) led to a significant attenuation in ear blood flow after niacin to  $143\% \pm 15\%$  of baseline flow, whereas TPAU alone at the same dose was ineffective in altering blood flow in the absence of niacin treatment (Fig. 2, open squares). Reduction in ear blood flow by TPAU was dose dependent and significant at 0.4 and 0.1 mg/kg (Fig. 3).

The sEH TPAU (0.4 mg/kg) was as effective as aspirin or the selective COX-2 inhibitor celecoxib in reducing vasodilation after niacin treatment (TPAU  $159 \pm 30$  vs. aspirin  $205 \pm 44$  and celecoxib  $222\% \pm 64\%$  of baseline blood flow), although at a 10-fold lower dose (Fig. 4). The commercial raf-1 kinase inhibitor sorafenib ( $185\% \pm 39\%$  of baseline)<sup>19</sup> and t-AUCB ( $149\% \pm 26\%$  of baseline), both potent sEH inhibitors with structures different than TPAU, similarly decreased niacin-induced flushing (Fig. 5). Taken




**FIGURE 2.** Pharmacologic inhibition of the sEH enzyme attenuated niacin-induced skin vasodilation. The sEH inhibitor TPAU 1 mg/kg (n = 4), when administered 45 minutes before niacin, significantly limited ear blood flow, whereas TPAU + vehicle alone (n = 3) did not alter skin blood flow (open squares) in the absence of niacin (see **Supplemental Digital Content 1**, <http://links.lww.com/JCVP/A76>, for the structure of TPAU). Abbreviations same as Figure 1; data for WT mice given niacin are data from Figure 1. WT, wild type.


together, these data demonstrate that inhibition of sEH, either through genetic deletion of the sEH gene or through the use of structurally distinct sEH inhibitors, was responsible for the observed biologic effects.



**FIGURE 1.** Targeted deletion of the sEH gene attenuated niacin-induced skin vasodilation. Ear blood flow (expressed as a percentage of baseline flow) in WT (n = 8) and KO (sEH -/-, n = 6) mice given niacin. The normal niacin response was characterized by a 3- to 5-fold increase in ear blood flow, which was significantly attenuated in the sEH KO mice. %BF (TPU), percent increase in ear blood flow (tissue perfusion units). WT, wild type.




**FIGURE 3.** Ear blood flow 4 minutes after niacin administration in mice pretreated with increasing doses of TPAU (n = 4 for niacin without TPAU and niacin with 0.04 and 0.4 mg/kg TPAU, n = 3 for niacin with 0.01 and 1 mg/kg TPAU, and TPAU without niacin) displayed a dose-dependent reduction in vasodilation. Due to potential niacin “tolerance,” each experimental animal was used only once and at 1 dose of niacin (30 mg/kg) and sEH. Abbreviations same as Figure 1. \* $P < 0.05$ .




**FIGURE 4.** Time response of skin vasodilation of TPAU compared with celecoxib and aspirin. Inhibition of sEH was more effective than inhibition of COX in reducing niacin-induced skin vasodilation. All compounds significantly inhibited the flushing response to niacin, although TPAU was effective at an order of magnitude lower concentration than the COX-1 and COX-2 selective inhibitors (niacin  $n = 6$ , all other groups  $n = 4$  per group). Abbreviations same as Figure 1; data for WT mice given niacin are data from Figure 1. WT, wild type.

### Niacin Reduces EETs, Whereas Inhibition of sEH Restores EETs to Baseline Values

Ear tissue and plasma levels of EETs and DHETs were measured to validate the efficacy of pharmacologic inhibition of sEH on cyp450 metabolism. Niacin alone reduced the levels of EETs in plasma from  $28.2 \pm 2.9$  to  $16.3 \pm 1.2$  nM, whereas pretreatment with TPAU restored the levels of EETs to  $25.3 \pm 2.7$  nM. There was no corresponding change in the overall levels of DHETs, resulting in a niacin-induced reduction in the ratio of EETs to DHETs (Fig. 6). However, when examining individual epoxyeicosatrienoic and DHETs, it was noted that whereas niacin did not significantly affect the plasma level of 14,15 DHET, pretreatment with TPAU



**FIGURE 5.** Maximum ear blood flow 4 minutes after niacin administration in mice pretreated with 3 sEH inhibitors ( $n = 4$  per group except for sorafenib,  $n = 3$ ) compared with aspirin and celecoxib (\*ANOVA, all groups  $P < 0.05$  vs. niacin alone).



**FIGURE 6.** Levels of EETs and DHETs in ear tissue and plasma obtained at the same time as samples in Figures 6 and 7. Niacin significantly reduced EETs/DHETs in plasma; this effect was reversed by pretreatment with sEH. \* $P < 0.05$  niacin versus other groups.

significantly reduced 14,15 DHET (control:  $0.47 \pm 0.04$ ; niacin:  $0.39 \pm 0.06$ ; niacin + TPAU:  $0.16 \pm 0.01$  nM;  $P < 0.05$ ).


### Inhibition of sEH Does Not Decrease Vasodilation Elicited by Systemically Administered PGD<sub>2</sub>

One potential mechanism for the above observations is a reduced vasodilator response to PGD<sub>2</sub> by a direct effect of sEH inhibition. To investigate whether sEH inhibition reduced the vasodilator response to PGD<sub>2</sub>, the flushing response in mice produced by direct SC administration of PGD<sub>2</sub> at 1 mg/kg was measured with and without sEH inhibition (TPAU 1 mg/kg) and in sEH  $-/-$  mice. Vasodilation produced by niacin and PGD<sub>2</sub> displayed different qualitative properties (Fig. 7), with systemic administration of PGD<sub>2</sub> resulting in a steady increase in cutaneous vasodilation over the course of 30 minutes, whereas niacin increased cutaneous vasodilation rapidly and transiently (Fig. 1). The vasodilatory response to PGD<sub>2</sub> was not significantly altered in either the sEH  $-/-$  mice or in the mice pretreated with TPAU. These findings indicate that sEH, either pharmacologic or genetic, did not inhibit the vasodilator response to exogenous PGD<sub>2</sub>.

## DISCUSSION

These experiments demonstrate that niacin-induced flushing can be blocked by genetic or pharmacologic inhibition of the key enzyme in the cyp450 system of ARA metabolism, sEH. The effect is qualitatively similar to that of conventional agents inhibiting the COX pathway, albeit at 1/10 the concentration. Furthermore, the effect of sEH inhibition does not appear to involve the response to PGD<sub>2</sub>.

Niacin has the potential to significantly improve cardiovascular health in a considerable number of patients.



**FIGURE 7.** Ear blood flow induced by SC administration of PGD<sub>2</sub> (1 mg/kg) was not different in KO (sEH  $-/-$ , n = 4) mice or WT mice pretreated with the sEH inhibitor TPAU (1 mg/kg, n = 4) compared with wild type without TPAU pretreatment (n = 5). This profile of vasodilation was markedly different than that produced by niacin (Fig. 1). WT, wild type.

However, the major obstacle in its widespread use, cutaneous vasodilation, is not well addressed. This side effect is serious enough that niacin is now used as an aversive agent against opioid abuse in formulations of opioids.<sup>20</sup> One major hypothesis of its mechanism of action in inducing flushing is a cascade of events initiated by the binding of niacin to the GPR109A receptor, leading to activation of phospholipase A<sub>2</sub> and the formation of ARA, which then is oxidized to the vasodilatory prostanoids PGD<sub>2</sub> and PGE<sub>2</sub>. Even though the GPR109A receptor is widely expressed, the flushing response is thought to be mediated by the epidermal Langerhans cells and is therefore a local rather than systemic event. PGD<sub>2</sub> and PGE<sub>2</sub> produced in the epidermal Langerhans cells can act on subepidermal blood vessels to induce vasodilation via activation of prostaglandin DP1 and EP<sub>2</sub>/EP<sub>4</sub> receptors.<sup>2</sup> Recently, experiments by Hanson et al<sup>21</sup> demonstrated that the biphasic vasodilatory response to niacin (as seen in Fig. 1) could be explained by an initial release of PGD<sub>2</sub> in Langerhans cells, whereas the second phase was due to PGE<sub>2</sub> release in keratinocytes.

The hypothesis that PGD<sub>2</sub> mediates the flushing effect of niacin has been tested in the murine model of niacin-induced flushing<sup>8</sup> and in cell lines and tissue samples.<sup>9</sup> In support of this hypothesis, interventions that reduce prostanoid synthesis from ARA (eg, aspirin), and specific antagonists of the DP1 receptor, partially reduced vasodilation in mice and symptomatic flushing in man.<sup>8,10</sup> However, because these agents are not fully effective, other mechanisms of niacin-induced flushing, including niacin-induced release of serotonin from platelets, have also been suggested.<sup>22</sup>

The efficacy of sEH inhibition in the current experiments was demonstrated using sEH KO mice and pharmacologic

inhibition with 3 structurally different inhibitors. Conventional predominantly COX-1 and COX-2 inhibitors, for example, aspirin and celecoxib, were also effective in reducing flushing in mice (Fig. 5), although equivalent efficacy of the sEH inhibitor TPAU was seen at one-tenth the dose of the COX inhibitors. Although the use of sEH inhibitors has not been reported in this flushing model, the data with COX inhibition are consistent with prior animal studies,<sup>8</sup> thereby validating the use of this model.

The efficacy of sEH inhibition was evaluated using measurements of EETs and DHETs. A novel finding in this study was the significant reduction of EETs, and the ratio of EETs to DHETs, by niacin alone. Coupled with previous reports of increased prostaglandins in response to niacin,<sup>22</sup> these data support a shift of ARA metabolism from the cyp450 pathway to the COX pathway and, potentially, the lipoxygenase pathway.<sup>23</sup> As expected, inhibition of sEH increased EETs,<sup>12</sup> resulting in normalization of the ratio of EETs to DHETs. Because inhibitors of sEH not only stabilize the levels of EETs but also modulate the metabolism of ARA by shifting flow from the COX branch toward the other 2 branches,<sup>13</sup> it is possible that the mechanism of sEH inhibition was through reduced prostaglandin production and/or a change in lipoxygenase products.<sup>24</sup>

Both EETs and DHETs have been shown to be direct nonendothelial dependent vasodilatory compounds, including coronary arteries<sup>25</sup> and capillary beds.<sup>26</sup> It is unclear why niacin, which results in vasodilation, would result in lower levels of EETs, both in absolute levels and when compared to DHETs. However, it is possible that the mechanisms of niacin-induced vasodilation are independent of either EETs and DHETs or that the relative abundance of DHETs relative to EETs in the skin vascular bed after niacin results in vasodilation. An intriguing possibility is that because various EETs and DHETs may have different vasodilatory effects in different vascular beds,<sup>27</sup> the reduction in 14,15 DHET may be contributory to the results seen with TPAU. Confirmation of this would clearly require further experimentation.

## Limitations

These experiments were performed in an established mouse model of niacin-induced flushing<sup>8</sup> using male animals and SC administration of both niacin and the sEH inhibitors. Therefore, it is unknown whether similar effects would be observed in other animal species (eg, humans) with oral dosing. However, the successful translation of animal results using laropiprant to humans suggests that this model is appropriate to determine efficacy.<sup>8,28</sup> In addition, although it is clear that sEH inhibition reduced flushing with niacin, the mechanism of flushing reduction is unknown. However, the preserved response to exogenous PGD<sub>2</sub> indicates that the effect of sEH inhibition is upstream of the DP1 receptor.

## CONCLUSIONS

Overall, our findings indicate that niacin-induced flushing is, in part, dependent on the cyp450 pathway of ARA metabolism because inhibition of this pathway, either using sEH KO mice or pharmacologic inhibition of sEH with

structurally different sEH inhibitors, blocks the flushing response. Furthermore, the inhibitory effect of sEH inhibition is not due to decreased responsiveness to PGD<sub>2</sub>. Therefore, sEH inhibition may limit niacin-induced flushing by countering the shift of ARA metabolism from the COX pathway to others such as the lipoxygenase pathways.<sup>24</sup> These findings strongly argue that inhibition of sEH may be a new therapeutic strategy to attenuate the major side effect of niacin therapy, thereby increasing patient compliance.

## REFERENCES

- Canner PL, Berge KG, Wenger NK, et al. Fifteen year mortality in Coronary Drug Project patients: long-term benefit with niacin. *J Am Coll Cardiol*. 1986;8:1245–1255.
- Gille A, Bodor ET, Ahmed K, et al. Nicotinic acid: pharmacological effects and mechanisms of action. *Annu Rev Pharmacol Toxicol*. 2008; 48:79–106.
- Berge KG, Canner PL. Coronary drug project: experience with niacin. Coronary Drug Project Research Group. *Eur J Clin Pharmacol*. 1991;40 (suppl 1):S49–S51.
- Davidson MH. Niacin use and cutaneous flushing: mechanisms and strategies for prevention. *Am J Cardiol*. 2008;101:14B–19B.
- Kamal-Bahl S, Watson DJ, Ambegaonkar BM. Patients' experiences of niacin-induced flushing in clinical practice: a structured telephone interview. *Clin Ther*. 2009;31:130–140.
- Cefali EA, Simmons PD, Stanek EJ, et al. Aspirin reduces cutaneous flushing after administration of an optimized extended-release niacin formulation. *Int J Clin Pharmacol Ther*. 2007;45:78–88.
- Hussein AA, Nicholls SJ. Critical appraisal of laropiprant and extended-release niacin combination in the management of mixed dyslipidemias and primary hypercholesterolemia. *Ther Clin Risk Manag*. 2010;6:183–190.
- Cheng K, Wu TJ, Wu KK, et al. Antagonism of the prostaglandin D2 receptor 1 suppresses nicotinic acid-induced vasodilation in mice and humans. *Proc Natl Acad Sci U S A*. 2006;103:6682–6687.
- Maciejewski-Lenoir D, Richman JG, Hakak Y, et al. Langerhans cells release prostaglandin D2 in response to nicotinic acid. *J Invest Dermatol*. 2006;126:2637–2646.
- Paolini JF, Bays HE, Ballantyne CM, et al. Extended-release niacin/laropiprant: reducing niacin-induced flushing to better realize the benefit of niacin in improving cardiovascular risk factors. *Cardiol Clin*. 2008;26: 547–560.
- Spector AA, Norris AW. Action of epoxyeicosatrienoic acids on cellular function. *Am J Physiol Cell Physiol*. 2007;292:C996–C1012.
- Spector AA. Arachidonic acid cytochrome P450 epoxygenase pathway. *J Lipid Res*. 2009;50(suppl):S52–S56.
- Liu JY, Yang J, Inceoglu B, et al. Inhibition of soluble epoxide hydrolase enhances the anti-inflammatory effects of aspirin and 5-lipoxygenase activation protein inhibitor in a murine model. *Biochem Pharmacol*. 2010;79:880–887.
- Iliff JJ, Wang R, Zeldin DC, et al. Epoxyeicosanoids as mediators of neurogenic vasodilation in cerebral vessels. *Am J Physiol Heart Circ Physiol*. 2009;296:H1352–H1363.
- Luria A, Weldon SM, Kabacnel AK, et al. Compensatory mechanism for homeostatic blood pressure regulation in Ephx2 gene-disrupted mice. *J Biol Chem*. 2007;282:2891–2898.
- Jones PD, Tsai HJ, Do ZN, et al. Synthesis and SAR of conformationally restricted inhibitors of soluble epoxide hydrolase. *Bioorg Med Chem Lett*. 2006;16:5212–5216.
- Morisseau C, Goodrow MH, Newman JW, et al. Structural refinement of inhibitors of urea-based soluble epoxide hydrolases. *Biochem Pharmacol*. 2002;63:1599–1608.
- Yang J, Schmelzer K, Georgi K, et al. Quantitative profiling method for oxylipin metabolome by liquid chromatography electrospray ionization tandem mass spectrometry. *Anal Chem*. 2009;81:8085–8093.
- Liu JY, Park SH, Morisseau C, et al. Sorafenib has soluble epoxide hydrolase inhibitory activity, which contributes to its effect profile in vivo. *Mol Cancer Ther*. 2009;8:2193–2203.
- Raffa RB, Pergolizzi JV Jr. Opioid formulations designed to resist/deter abuse. *Drugs*. 2010;70:1657–1675.
- Hanson J, Gille A, Zwykiel S, et al. Nicotinic acid- and monomethyl fumarate-induced flushing involves GPR109A expressed by keratinocytes and COX-2-dependent prostanoid formation in mice. *J Clin Invest*. 2010;120:2910–2919.
- Papalioidis D, Boucher W, Kempuraj D, et al. Niacin-induced “flush” involves release of prostaglandin D2 from mast cells and serotonin from platelets: evidence from human cells in vitro and an animal model. *J Pharmacol Exp Ther*. 2008;327:665–672.
- Saareks V, Mucha I, Sievi E, et al. Nicotinic acid and pyridoxine modulate arachidonic acid metabolism in vitro and ex vivo in man. *Pharmacol Toxicol*. 1999;84:274–280.
- Chawengsub Y, Aggarwal NT, Nithipatikom K, et al. Identification of 15-hydroxy-11,12-epoxyeicosatrienoic acid as a vasoactive 15-lipoxygenase metabolite in rabbit aorta. *Am J Physiol Heart Circ Physiol*. 2008; 294:H1348–H1356.
- Oltman CL, Weintraub NL, VanRollins M, et al. Epoxyeicosatrienoic acids and dihydroxyeicosatrienoic acids are potent vasodilators in the canine coronary microcirculation. *Circ Res*. 1998;83:932–939.
- Harder DR, Campbell WB, Roman RJ. Role of cytochrome P-450 enzymes and metabolites of arachidonic acid in the control of vascular tone. *J Vasc Res*. 1995;32:79–92.
- Larsen BT, Miura H, Hatoum OA, et al. Epoxyeicosatrienoic and dihydroxyeicosatrienoic acids dilate human coronary arterioles via BK(Ca) channels: implications for soluble epoxide hydrolase inhibition. *Am J Physiol Heart Circ Physiol*. 2006;290:H491–H499.
- Paolini JF, Mitchel YB, Reyes R, et al. Effects of laropiprant on nicotinic acid-induced flushing in patients with dyslipidemia. *Am J Cardiol*. 2008; 101:625–630.