



Lipoxin Generation Is Related to Soluble Epoxide Hydrolase Activity in Severe Asthma

Emiko Ono¹, Stefanie Dutile¹, Shamsah Kazani¹, Michael E. Wechsler¹, Jun Yang², Bruce D. Hammock², David Nobuhiro Doua¹, Yacine Tabet³, Rayan Khaddaj-Mallat³, Marco Sirois⁴, Chantal Sirois⁴, Edmond Rizcallah⁵, Éric Rousseau³, Richard Martin⁶, E. Rand Sutherland⁶, Mario Castro⁷, Nizar N. Jarjour⁸, Elliot Israel¹, Bruce D. Levy¹, and National Heart, Lung, and Blood Institute's Asthma Clinical Research Network*

¹Pulmonary and Critical Care Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts; ²Department of Entomology and Comprehensive Cancer Center, University of California at Davis, Davis, California; and ³Le Bilarium, Department of Physiology and Biophysics, ⁴Service of Thoracic Surgery, and ⁵Department of Pathology, Faculty of Medicine and Health Sciences, Université de Sherbrooke, Sherbrooke, Quebec, Canada; ⁶Department of Medicine, National Jewish Health, Denver, Colorado; ⁷Pulmonary and Critical Care Medicine, Washington University in St. Louis, St. Louis, Missouri; and ⁸Allergy, Pulmonary and Critical Care Medicine, University of Wisconsin, Madison, Wisconsin

Abstract

Rationale: Severe asthma is characterized by airway inflammatory responses associated with aberrant metabolism of arachidonic acid. Lipoxins (LX) are arachidonate-derived pro-resolving mediators that are decreased in severe asthma, yet mechanisms for defective LX biosynthesis and a means to increase LXs in severe asthma remain to be established.

Objectives: To determine if oxidative stress and soluble epoxide hydrolase (sEH) activity are linked to decreased LX biosynthesis in severe asthma.

Methods: Aliquots of blood, sputum, and bronchoalveolar lavage fluid were obtained from asthma subjects for mediator determination. Select samples were exposed to *t*-butyl-hydroperoxide or sEH inhibitor (sEHI) before activation. Peripheral blood leukocyte-platelet aggregates were monitored by flow cytometry, and bronchial contraction was determined with cytokine-treated human lung sections.

Measurements and Main Results: 8-Isoprostane levels in sputum supernatants were inversely related to LXA₄ in severe asthma ($r = -0.55$; $P = 0.03$) and *t*-butyl-hydroperoxide decreased LXA₄ and 15-epi-LXA₄

biosynthesis by peripheral blood leukocytes. LXA₄ and 15-epi-LXA₄ levels were inversely related to sEH activity in sputum supernatants and sEHs significantly increased 14,15-epoxy-eicosatrienoic acid and 15-epi-LXA₄ generation by severe asthma whole blood and bronchoalveolar lavage fluid cells. The abundance of peripheral blood leukocyte-platelet aggregates was related to asthma severity. In a concentration-dependent manner, LXs significantly inhibited platelet-activating factor-induced increases in leukocyte-platelet aggregates (70.8% inhibition [LXA₄ 100 nM], 78.3% inhibition [15-epi-LXA₄ 100 nM]) and 15-epi-LXA₄ markedly inhibited tumor necrosis factor- α -induced increases in bronchial contraction.

Conclusions: LX levels were decreased by oxidative stress and sEH activity. Inhibitors of sEH increased LXs that mediated antiphlogistic actions, suggesting a new therapeutic approach for severe asthma.

Clinical trial registered with www.clinicaltrials.gov (NCT 00595114).

Keywords: resolution; asthma; pro-resolving mediators; inflammation

Severe asthma (SA) is distinguished from milder variants by reduced corticosteroid responsiveness (1, 2) that derives in part

from increased airway oxidative stress (3). No matter the cause of this heterogeneous condition (4), most patients with SA

experience excess, chronic airway inflammation with defective resolution (5, 6).

(Received in original form March 21, 2014; accepted in final form August 21, 2014)

*A complete list of members may be found before the beginning of the REFERENCES.

Supported in part by National Institutes of Health grants HL107166, HL090927, HL109172, HL102225, ES002710, ES004699, and DK097154 and CIHR grant MOP-111112. B.D.H. is a George and Judy Marcus Fellow of the American Asthma Society and É.R. is a member of the Centre de Recherche Clinique E. LeBel, CHUS, Sherbrooke.

Author Contributions: Conception and experimental design, E.O., É.R., E.I., and B.D.L. Sample and data acquisition, all authors. Manuscript preparation, E.O., M.E.W., J.Y., B.D.H., Y.T., É.R., E.I., and B.D.L.

Correspondence and requests for reprints should be addressed to Bruce D. Levy, M.D., Pulmonary and Critical Care Medicine, Brigham and Women's Hospital/Harvard Medical School, 77 Avenue Louis Pasteur, Boston, MA 02115. E-mail: blevy@partners.org

This article has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org

Am J Respir Crit Care Med Vol 190, Iss 8, pp 886–897, Oct 15, 2014

Copyright © 2014 by the American Thoracic Society

Originally Published in Press as DOI: 10.1164/rccm.201403-0544OC on August 27, 2014

Internet address: www.atsjournals.org

At a Glance Commentary

Science Knowledge on the

Subject: In severe asthma, the levels of lipoxins (LXs) are decreased by post-transcriptional mechanisms that have yet to be defined. Because LXs are pro-resolving mediators, defective LX generation may underlie the chronic inflammatory airway changes in severe asthma. Because LX formation occurs via transcellular biosynthesis during cell–cell interactions, these biosynthetic pathways are potentially vulnerable to oxidative stress.

What This Study Adds to the

Field: In severe asthma, LX generation in the airway was inversely related to oxidative stress and soluble epoxide hydrolase activity. Highly selective inhibitors of soluble epoxide hydrolase increased 14,15-epoxy-eicosatrienoic acid and 15-epi-LXA₄ generation by severe asthma whole blood and bronchoalveolar lavage cells. 15-epi-LXA₄ decreased leukocyte–platelet aggregates and both 15-epi-LXA₄ and 14,15-epoxy-eicosatrienoic acid decreased bronchial contraction.

The resolution of inflammation is an active process governed by specialized pro-resolving mediators and cellular events (7). In model systems, pro-resolving mediators can potentially decrease further recruitment of leukocytes to inflamed tissues and promote a return of the tissue to homeostasis (8). The genus of endogenous pro-resolving mediators includes lipoxins (LXs), which are enzymatically derived from arachidonic acid during cell–cell interactions in inflammation (9). The multistep enzymatic process of LX biosynthesis with transcellular exchange of biosynthetic intermediates may increase its vulnerability to corruption by nonenzymatic, oxidative attack on the arachidonate backbone in the presence of increased oxidative stress.

LXs are generated in asthmatic airways (10); however, LX generation in SA is defective (11–13). LXs can regulate allergic airways responses, including airway inflammation, mucus metaplasia, and hyperresponsiveness to methacholine (MCh) (14). At a cellular level, LXs inhibit

eosinophil trafficking (14) and neutrophil chemotaxis, transendothelial and transepithelial migration, generation of superoxide anions, and degranulation of neutrophil azurophilic granules (7). In addition, inhaled LXA₄ blocks leukotriene (LT) C₄ (LTC₄)-mediated bronchoprovocation of subjects with asthma (15) and a LXA₄ stable analog markedly decreases infantile eczema (16). SA is a uniquely human disease and important species-specific differences are present between LX biosynthetic enzymes, emphasizing the importance of human translational research in this condition.

Here, we determine the relationship between LX levels and oxidative stress, identify a post-transcriptional mechanism to increase LX biosynthesis in SA, and uncover protective actions for LXs on two important pathologic features of SA, namely platelet–leukocyte interactions for lung leukocyte recruitment and proinflammatory cytokine induced *ex vivo* bronchial reactivity.

Methods

See the online supplement for detailed methods.

Participants

Aliquots of materials were collected from a random subset of individuals enrolled in the Macrolides In Asthma (MIA; NCT 00318708) trial or the cKit Inhibition in Severe Asthma (KIA; NCT 01097694) trial for this ancillary mechanistic study. Asthma severity was graded nonsevere asthma (NSA) or SA based on criteria developed by the NHLBI Severe Asthma Research Program (4). The Brigham and Women's Hospital human subjects institutional review board approved the study, and all subjects provided written consent in accordance with the Declaration of Helsinki.

Sample Collection

Peripheral venous blood was collected in heparinized tubes and used immediately. Induced sputum was prepared as in (13) with supernatants stored at -80°C for later analysis. Bronchoalveolar lavage fluids (BALF) were collected as in (17) with cell-free supernatants stored at -80°C as methanolic extracts (1:1, vol/vol BALF/methanol). All samples were collected from volunteer subjects before the initiation of study medication or placebo. Prostaglandin B₂ was added as an internal control.

Effects of *t*-Butyl-Hydroperoxide and Soluble Epoxide Hydrolase Inhibitors

Partially purified peripheral blood leukocytes were warmed (5 min, 37°C) and exposed (15 min, 37°C) to *t*-butyl-hydroperoxide (*t*-BuOOH; 2 mM) or vehicle before incubation with A23187 (50 μM) or vehicle (0.1% ethanol) for 15 minutes (37°C). Incubations were stopped with 5 vols of iced methanol and stored at -80°C . In separate experiments, peripheral venous blood (700 μl per incubation) was warmed (5 min, 37°C) and exposed (10 min, 37°C) to either a soluble epoxide hydrolase inhibitor (sEHI) (500 nM; 12-[3-adamantan-1-yl-ureido] dodecanoic acid [AUDA], 1-[1-acetylpiperidin-4-yl]-3-adamantanyleurea, 1-trifluoromethoxyphenyl-3-[1-propionylpiperidin-4-yl] urea, and *trans*-4-[4-[3-(4-trifluoromethoxy-phenyl)-ureido]-cyclohexyloxy]-benzoic acid [*t*-TUCB]; structures given in online supplement) or vehicle (0.1% dimethyl sulfoxide) before A23187 (50 μM , 15 min, 37°C). Incubations were stopped with 5 vols of iced methanol. BALF cells (1×10^5 cells per incubation) in phosphate-buffered saline were warmed (5 min, 37°C) and exposed (10 min, 37°C) to either a sEHI (500 nM) or vehicle before A23187 (5 μM , 15 min, 37°C). Incubations were stopped with 2 vols of iced methanol.

Extraction and Measurement of Lipid Mediators

Whole blood and BALF cell incubations were extracted as in (12, 17). Quantitation of LXA₄, 15-epi-LXA₄, LTB₄, 8-isoprostane (8-IP), 14,15-epoxy-eicosatrienoic acid (14,15-EET), and 14,15-dihydroxy-eicosatrienoic acid (14,15-DHET) was determined in parallel by ELISA (Neogen, Lansing, MI; Cayman Chemical, Ann Arbor, MI; Detroit R&D, Detroit, MI). LXA₄ levels were confirmed by liquid chromatography–mass spectrometry in select samples as in (18).

Flow Cytometry

Freshly obtained peripheral venous blood was obtained from subjects enrolling at the Boston site for the MIA or KIA trials or from healthy control (HC) subjects. Leukocyte–platelet aggregates (LPAs) were analyzed by flow cytometry as in (19). LPAs were identified as CD16⁺CD41⁺. To some aliquots of whole blood, samples were exposed to either LXA₄, 15-epi-LXA₄, 14,15-EET, 8-IP (Cayman Chemical), or

vehicle (0.1% ethanol) (5 min, 37°C) before activation with platelet-activating factor (PAF; 5 μ M) or vehicle (0.1% ethanol) (15 min, 37°C). To stimulate endogenous LX generation, samples of whole blood were first exposed to granulocyte-macrophage colony-stimulating factor (200 pM, 90 min, 37°C) before sEHI or vehicle (10 min, 37°C) and followed by activation with PAF (15 min, 37°C). At least 20,000 CD41⁺ cells were recorded for each sample on a FACS Canto II (BD Biosciences, San Jose, CA) and data were analyzed using FloJo software (Tree Star, Ashland, OR).

Human Lung Tissue Samples

The study was approved by the Université de Sherbrooke's Ethics Committee (protocol number 05-088-S1-R2). Human lung tissues were obtained from 10 patients undergoing resection and bronchi were dissected and cultured for 48 hours with 10 ng/ml tumor necrosis factor (TNF)- α (R&D Systems, Minneapolis, MN) in the presence of 100 nM 15-epi-LXA₄, 14,15-EET, or vehicle.

Isometric Tension Measurements

Mechanical tension measurements were performed using an isolated organ bath system (Radnoti Glass Tech., Monrovia, CA) as previously described (20). Briefly, bronchial rings were mounted in baths filled with 6 ml of Krebs solution, at pH 7.4 (37°C, 5% CO₂). A basal tension of 0.8 g was applied to each bronchi. Passive and active tensions were assessed using FT03 Grass transducer systems coupled to Polyview software (Grass-Astro-Med Inc., West Warwick, RI) to allow data acquisition and analysis.

Statistical Analysis

Samples were deidentified before analysis. Analysis was based on the Mann-Whitney test, the Kruskal-Wallis analysis of variance followed by the Mann-Whitney test with Bonferroni correction to compare the ranks of continuous variables across the levels of a categorical variable with more than two groups, the Spearman rank correlation test to compare the ranks of two continuous variables, and chi-square tests and Fisher exact test for comparisons of categorical variables. Data are presented as the mean \pm SD or SEM where indicated. *P* less than 0.05 was regarded as statistically significant. Prism (GraphPad Software, La Jolla, CA) or Sigma Plot 12.0 (SPSS-Science, Chicago, IL) were used to manage and analyze the data.

Results

Subject Characteristics

As an ancillary study, samples were obtained from a random subset of subjects with asthma participating in either the MIA or KIA trials. Using criteria developed by the Severe Asthma Research Program (4), participants could be separated into cohorts of SA and NSA. The clinical profiles of the enrolled subjects with NSA and SA whose materials were studied here are provided in Table 1. Relative to NSA, subjects with SA used significantly more inhaled corticosteroids and long-acting bronchodilators with higher Asthma Control Questionnaire scores. Healthy subjects were also recruited as control subjects (HC) (*see* online supplement for detailed methods).

Lipid Mediators and 8-IP in NSA Plasma and Sputum

To determine if the abundance of lipoygenase-derived (LXs and LTs) and nonenzymatically derived (IPs) eicosanoids differ by anatomic compartment, levels of representative members of these families (i.e., LXA₄, LTB₄, and 8-IP) were measured in samples of plasma and sputum from subjects with NSA (Figure 1). LXA₄, LTB₄, and 8-IP were detected in all samples (Figures 1A and 1B). Plasma levels of LXA₄ (mean \pm SD, 463.3 \pm 152.6 pg/ml) were significantly higher than LTB₄ (mean \pm SD, 16.8 \pm 18.8 pg/ml; *P* < 0.05) (Figure 1A), but not in sputum supernatants (Figure 1B). Substantial amounts of 8-IP were present in both

plasma (mean \pm SD, 254.0 \pm 125.6 pg/ml) and sputum (mean \pm SD, 292.4 \pm 51.0 pg/ml) (Figures 1A and 1B). There was no significant relationship between levels of LXA₄, LTB₄, and 8-IP in plasma and sputum (Figure 1C), suggesting important anatomic differences in arachidonic acid availability and metabolism. The levels of these mediators were not significantly related to the clinical parameters listed in Table 1.

LXA₄ Generation Is Inversely Related to Oxidative Stress in the SA Airway

To determine if there was a relationship between LX generation and oxidative stress, levels of LXA₄ and 8-IP, a sensitive biomarker for oxidative stress, were measured in samples of plasma and sputum. Of interest, there was a significant positive correlation between LXA₄ and 8-IP in NSA plasma (*r* = 0.68; *P* = 0.002) (Figure 2A), suggesting that plasma levels of these compounds were related to arachidonic acid availability. Unlike plasma, there was a negative relationship and no significant correlation between LXA₄ and 8-IP levels in NSA sputum supernatants (*r* = -0.40; *P* = 0.28) (Figure 2B).

In sharp contrast with NSA, a significant inverse relationship between LXA₄ and 8-IP levels in SA sputum supernatants was present (*r* = -0.55; *P* = 0.03) (Figure 2C), indicative of an inverse relationship between LX generation and oxidative stress in the airway, in particular in SA. To this end, the mean levels of 8-IP were higher in SA sputum supernatants

Table 1. Subject Characteristics

	Mild Asthma	Severe Asthma
Number of subjects	24	19
Clinical data		
Age, yr	37 \pm 10 (22–58)	40 \pm 11 (20–52)
Sex, M/F	7/17	10/9
Race, % white	37.5	57.9
Ethnicity, % Hispanic	20.8	15.8
Inhaled steroids dose, μ g LABA/Tio, %	166 \pm 72 (0–320) 0	922 \pm 320* (427–1807) 100*
Lung function		
ACQ score	1.5 \pm 0.65 (0.43–3.4)	2.6 \pm 0.76* (2.0–4.7)
FEV ₁ , L	2.4 \pm 0.68 (1.1–3.9)	2.4 \pm 0.74 (1.3–4.1)
FEV ₁ , % predicted	77 \pm 18 (38–106)	72 \pm 19 (40–101)

Definition of abbreviations: ACQ = Asthma Control Questionnaire; LABA = long-acting β agonist. Results are expressed as mean \pm SD (range).

**P* < 0.05 when compared with subjects with mild asthma.




Figure 1. The relationship between lipoxin (LX) A₄, leukotriene (LT) B₄, and 8-isoprostane levels differs by anatomic compartment in asthma. Samples were obtained from a subset of subjects with nonsevere asthma enrolled in the Macrolides In Asthma trial. (A) Plasma (n = 17) and (B) sputum supernatants (n = 9) were extracted and levels of LXA₄, LTB₄, and 8-isoprostane were determined by ELISA (see METHODS). Results are expressed as individual values. The mean ± SEM are indicated by overlay in (A) plasma and (B) sputum. (C) The concentrations of LXA₄ (circles), LTB₄ (squares), and 8-isoprostane (triangles).

(mean ± SD, 421.8 ± 195.4 pg/ml) than NSA sputum supernatants (mean ± SD, 292.4 ± 51.0 pg/ml), but these differences did not reach statistical significance at this sample size (Figure 2D). To determine if LX biosynthesis was susceptible to oxidative stress, leukocytes from healthy subjects were partially purified from peripheral

blood and activated with A23187 in the presence or absence of *t*-BuOOH. Both LXA₄ and 15-epi-LXA₄ biosynthesis increased with A23187, but LX production was significantly inhibited by *t*-BuOOH (Figures 2E and 2F). Together, these findings are consistent with disruption of LX biosynthesis by oxidative stress.

Relationship between LXA₄ Levels and sEH Activity

sEH expression is increased by oxidative stress (21), abundant in asthmatic airways (22), and its activity can influence LX generation by human cells *in vitro* and in rodent lung *in vivo* (18, 23), so we next determined if there was a relationship between LXA₄ levels and sEH activity in the sputum supernatants from subjects with asthma (Figure 3). Because 14,15-EET is a prominent cytochrome P-450-derived eicosanoid in the lung that is converted by sEH to 14,15-DHET (24), the product/precursor ratio of 14,15-DHET to 14,15-EET was used as an index for sEH activity. Significant inverse relationships were identified between the ratio of 14,15-DHET to 14,15-EET and LXA₄ levels in both NSA (r = -0.81; P = 0.01) (Figure 3A) and SA sputum supernatants (r = -0.72; P = 0.002) (Figure 3B), indicative of a relationship between increased sEH activity and decreased LX levels. No significant correlations between ratio of 14,15-DHET to 14,15-EET and 8-IP were identified in these samples (data not shown).

sEH Decreases the Ratio of 14,15-DHET to 14,15-EET

In view of this inverse relationship between sEH activity and LXA₄ levels, we first tested the capacity of four potent and selective sEHIs (Table 2) to block sEH activity in samples from subjects with SA (Figure 4). In activated whole blood and BAL cells, all four sEHIs significantly decreased sEH activity relative to control incubations (P < 0.05) (Figures 4A and 4B). In addition to immunologic detection by ELISA, the identification of 14,15-EET and 14,15-DHET, and the impact of the sEHI on their formation was confirmed by physical methods with liquid chromatography–mass spectrometry (Figures 4C–4F) (see METHODS).

sEHI Increases 15-Epi-LXA₄ Biosynthesis

To determine if the sEHIs could increase LX generation, materials from the same

incubations in Figure 4 were also analyzed by ELISA for 15-epi-LXA₄. In the presence of A23187, 15-epi-LXA₄ was detectable in the incubations with whole blood (mean ± SD, 78.9 ± 30.9 pg/ml) and BAL cells (mean ± SD, 110.7 ± 65.2 pg/ml). There was a significant increase in 15-epi-LXA₄ levels with AUDA and *t*-TUCB in SA whole blood (Figure 5A) and with AUDA, 1-(1-acetylpiperidin-4-yl)-3-adamantanylurea, and *t*-TUCB with BALF cells (Figure 5B). The most active sEHs in these samples (Figures 4A and 4B) were generally also the most potent inducers of 15-epi-LXA₄ generation (Figures 5A and 5B). Similar to the relationship in SA and NSA sputum supernatants between endogenous LXA₄ and the 14,15-DHET/14,15-EET ratio (Figures 3A and 3B), there was also a significant inverse relationship between 15-epi-LXA₄ and the 14,15-DHET/14,15-DHET ratio with sEHs (Figure 5C).

Leukocyte–Platelet Aggregates Are Increased in SA and Regulated by LXs

Because leukocyte–platelet interactions are present in asthmatic blood (25) and involved in lung leukocyte recruitment (26, 27), we next determined if these heterotypic aggregates were related to asthma severity and regulated by LXs and sEH activity (Figure 6). LPAs were quantitated by determining the percent of granulocytes (identified by forward and side scatter, confirmed by CD16) that also expressed the platelet-specific marker CD41a (Figure 6A). The percentage was increased in SA relative to HC and could be increased even further by whole blood incubation with PAF (5 μM), a proinflammatory agonist for both leukocytes and platelets (Figures 6A and 6B). The percentage of LPAs was linked to asthma severity in the presence and absence of PAF (Figure 6B). LXA₄ (1–100 nM) inhibited PAF-induced LPAs in a concentration-dependent manner, but there was still segregation between the cohorts by asthma severity (Figure 6C). LXA₄ and 15-epi-LXA₄ (100 nM) were equipotent at inhibiting the PAF-induced formation of LPAs (mean percent inhibition; 70.8% for LXA₄, 78.3% for 15-epi-LXA₄), whereas 14,15-EET and 8-IP were inactive in this bioassay (Figure 6D). Incubation of whole blood from HC in the presence of *t*-TUCB and TPPU (chosen based on their ability to inhibit sEH and induce 15-epi-LXA₄) led to




Figure 2. Oxidative stress is inversely related to lipoxin (LX) levels. Samples were obtained from a subset of subjects with nonsevere asthma enrolled in the Macrolides In Asthma trial (same samples as in Figure 1) and subjects with severe asthma enrolled in the cKit Inhibition in Severe Asthma trial. Materials in (A) plasma and (B–D) sputum supernatants were extracted and levels of LXA₄ and 8-isoprostane, a sensitive indicator of oxidative stress, were determined by ELISA (see METHODS). Values for each individual sample are shown. Analysis was based on the Spearman rank correlation test (A, $r = 0.68$, $P = 0.002$; B, $r = -0.40$, $P = 0.28$; C, $r = -0.55$, $P = 0.03$). Black bar shows the 8-IP levels in sputum from SA subjects and white bar shows those from NSA subjects (D, $P = 0.17$). (E and F) To determine if LX production is disrupted by oxidative stress, peripheral blood leukocytes from healthy subjects were exposed to *t*-butyl-hydroperoxide (*t*-BuOOH) to purposely increase oxidative stress before activation of LX biosynthesis with A23187 (see METHODS). Results are expressed as the mean \pm SEM for $n \geq 3$. * $P < 0.05$ compared with vehicle/vehicle control, ** $P < 0.05$ compared with vehicle/A23187.

different levels of inhibition of PAF-induced cell-cell interactions (Figure 6E) that reflected their capacity to increase endogenous 15-epi-LXA₄ levels (Figure 6F).

Effect of 15-Epi-LXA₄ on TNF α -induced Bronchial Reactivity

In addition to LXs' potent regulation of LPAs for tissue inflammation, their direct actions on lung structural cells was next

determined with an *ex vivo* model of bronchial reactivity using human lung sections (Figure 7). Surgical specimens were obtained and exposed (48 h) to 15-epi-LXA₄ (100 nM), 14,15-EET (100 nM), or vehicle with TNF- α (10 ng/ml) to increase bronchial reactivity as in (28). Bronchi were challenged *in vitro* with a range of relevant pharmacologic agonists. Histamine (1 μ M) (Figure 7A) and U-46619 (30 nM, thromboxane receptor agonist) (Figure 7B) induced rapid increases in tension within seconds that reached a plateau by 5 minutes. With media alone, the mean tension for MCh (1 μ M), histamine (1 μ M), or U-46619 (30 nM) was 0.19 ± 0.03 g, 0.28 ± 0.04 g, and 0.34 ± 0.04 g, respectively (Figure 7C). TNF- α increased bronchial contractile responses to the same agents to 0.31 ± 0.04 g, 0.45 ± 0.07 g, and 0.55 ± 0.07 g, which represent significant increases in the active tension of 63%, 61%, and 62%, respectively, compared with media alone (Figure 7C). Exposure to 15-epi-LXA₄ (100 nM) markedly reversed the TNF- α -induced increases in mean tension on challenge with MCh to 0.17 ± 0.02 g, histamine to 0.27 ± 0.03 g, and U-46619 to 0.36 ± 0.04 g (Figure 7C). These findings with 15-epi-LXA₄ represent a significant reduction of 44.5, 38.6, and 34.4%, respectively, in TNF- α -mediated increases in bronchial contraction to a diverse range of stimuli. In addition to 15-epi-LXA₄, 14,15-EET also carried bioactivity in these assays. Direct exposure to 15-epi-LXA₄ reduced TNF- α -triggered responses by 113.5, 106.3, and 90.1% for MCh, histamine, and U-46619, respectively. For purposes of comparison with data included in our prior publication (28), exposure to 14,15-EET reduced TNF- α -induced bronchial contraction to these same agonists by 78.4, 73.4, and 85.2%, respectively (Figure 7D).

15-epi-LXA₄ can interact with multiple receptors in the airway, including as an agonist at ALX/FPR2 (29), so WRW4, a receptor antagonist for ALX/FPR2, was used to determine if the LX actions were ALX/FPR2-dependent. Paired recordings (Figures 7E and 7F) and quantitative analyses (Figure 7C) of bronchial responses to MCh revealed concentration-dependent bronchial relaxation by 15-epi-LXA₄ with partial inhibition by WRW4. Similar experiments were performed to assess the putative effects of WRW4 on the relaxing effects induced by 14,15-EET on MCh




Figure 3. Relationship between lipoxin (LX) A₄ levels and soluble epoxide hydrolase activity in asthma. Samples were obtained from (A) subjects with nonsevere asthma enrolled in the Macrolides In Asthma trial (same samples as in Figures 1 and 2) and (B) subjects with severe asthma enrolled in the cKit Inhibition in Severe Asthma trial (same samples as in Figure 2). Materials in sputum supernatants were extracted and levels of LXA₄, 14,15-epoxy-eicosatrienoic acid (14,15-EET), and 14,15-dihydroxy-eicosatrienoic acid (14,15-DHET) were determined by ELISA. The ratio of 14,15-DHET to 14,15-EET was calculated as an index of soluble epoxide hydrolase activity. Values for each individual sample are shown. Analysis was based on the Spearman rank correlation test (A, $r = -0.81$, $P = 0.02$; B, $r = -0.72$, $P = 0.002$).

precontracted human bronchi (see Figure E1 in the online supplement). 14,15-EET induced concentration-dependent

relaxation of mean MCh responses that were also partially inhibited by WRW4 (Figure 7F; see Figure E1). Together, these findings add direct tissue protective actions to 15-epi-LXA₄'s and 14,15-EET's antiasthmatic activities that are partially ALX/FPR2-dependent.

Discussion

The present findings provide evidence that oxidative stress and sEH activity are linked to decreased LX generation in SA. 8-IP levels were used as an arachidonic acid-based indicator of oxidative stress. This F2-IP is nonenzymatically synthesized by free radical-catalyzed lipid peroxidation (30). 8-IP is a sensitive biomarker of oxidative stress that is increased in asthma (30, 31) and related to disease activity (32). In contrast, LXs are generated enzymatically from arachidonic acid in a highly orchestrated, multistep biosynthesis that includes exchange of arachidonic acid and biosynthetic intermediates between cell types (33). These cell-cell interactions occur between leukocytes and platelets in the vasculature (33) and between leukocytes and mucosal epithelial cells in the airway (23). We hypothesized that the shuttling of polyunsaturated fatty acid intermediates between cells would increase their vulnerability in oxidative environments to nonenzymatic attack and subsequent disruption of LX biosynthesis. This susceptibility for LXs differs from LT biosynthesis, which often proceeds in single cells (9).

To address our hypothesis, we compared closely related eicosanoids (i.e., 8-IP, LXA₄, and LTB₄) in samples from patients with asthma. Levels of 8-IP were increased in SA sputum supernatants relative to NSA, and there was a significant inverse relationship between 8-IP and

LXA₄. Increased oxidative stress may lead to corticosteroid resistance, which is common in SA (3). LX levels are decreased in SA, yet corticosteroid dosing is not related to LX production in SA (17). Because the changes in LX levels here were linked to oxidative stress, indirectly reflected in the 8-IP levels, we purposefully induced oxidative stress in healthy leukocytes using t-BuOOH, which significantly disrupted LXA₄ and 15-epi-LXA₄ biosynthesis. In NSA sputum, there was also a trend toward an inverse relationship between LXA₄ and 8-IP, but it did not reach statistical significance and the slope was less marked. In sharp contrast, there was a positive correlation between 8-IP and LXA₄ in NSA plasma, which was more consistent with arachidonate availability as the limiting factor for plasma 8-IP, LTB₄, and LXA₄ and indicative of an important anatomic-specific influence of oxidative stress on LX biosynthesis.

LX levels in asthmatic sputum were also inversely related to sEH activity and were increased *in vitro* by sEHIs. sEH is expressed in asthmatic lung (22) and plays important roles in arachidonic acid metabolism (24). Expression of sEH is increased by oxidative stress (21). Cytochrome P-450 enzymes convert arachidonic acid to EETs, resulting in the production of four regioisomeric EETs: 5,6-EET, 8,9-EET, 11,12-EET, and 14,15-EET. Among these regioisomers, 14,15-EET is the most abundant in lung tissues (34). Based on protective actions of EETs and their sEH-mediated transformation to biologically inactive diols (DHETs), sEHIs have been developed (21, 35, 36). In rodents, inhibition of sEH decreases cigarette smoke-induced lung inflammation and increases LXA₄ levels (18). In addition, proinflammatory cytokine-exposed human airway epithelial cells can donate

Table 2. Four Specific Soluble Epoxide Hydrolase Inhibitors*

Compound	Abbreviation	Target Enzymes	IC ₅₀	Compound Name
EHI 700	AUDA	sEH and EH3	3 nM	12-(3-adamantan-1-yl-ureido) dodecanoic acid
EHI 1153	APAU	sEH	15 nM	1-(1-acetypiperidin-4-yl)-3-adamantanylurea
EHI 1770	TPPU	sEH	1 nM	1-trifluoromethoxyphenyl-3-(1-propionylpiperidin-4-yl) urea
EHI 1728	t-TUCB	sEH	2 nM	trans-4-{4-[3-(4-trifluoromethoxy-phenyl)-ureido]-cyclohexyloxy}-benzoic acid

Definition of abbreviations: EHI = epoxide hydrolase inhibitor; IC₅₀ = half maximal 50% inhibitory concentration; sEH = soluble epoxide hydrolase. *Structures of the sEHIs are shown in the online supplement.




Figure 4. Soluble epoxide hydrolase inhibitors decreased the ratio of 14,15-dihydroxy-eicosatrienoic acid (14,15-DHET) to 14,15-epoxy-eicosatrienoic acid (14,15-EET) in severe asthma. Samples of (A) whole blood and (B) bronchoalveolar lavage fluids were obtained from subjects with severe asthma and exposed to a soluble epoxide hydrolase inhibitor (sEHI) or vehicle before cell activation with A23187 (see METHODS). Materials from these incubations were extracted and a ratio of 14,15-DHET to 14,15-EET was determined as an indicator of sEH activity. Results are expressed as mean \pm SEM ($n \geq 6$). * $P < 0.05$ in comparison with samples activated by A23187 in the absence of sEHI by one-way analysis of variance. (C–F) Liquid chromatography–mass spectrometry (representative of $n = 3$). (C) Vehicle, (D) sEHI 700, (E) 14,15-EET, and (F) 14,15-DHET. APAU = 1-(1-acetypiperidin-4-yl)-3-adamantanyliurea; AUDA = 12-(3-adamantan-1-yl-ureido) dodecanoic acid; TPPU = 1-trifluoromethoxyphenyl-3-(1-propionylpiperidin-4-yl) urea; *t*-TUCB = *trans*-4-[4-[3-(4-trifluoromethoxy-phenyl)-ureido]-cyclohexyloxy]-benzoic acid.

14,15-EET to human neutrophils during coinubation for transcellular 15-epi-LXA₄ biosynthesis (23).

Because LX generation is deficient in SA, we determined the relationship between LXA₄ and sEH activity in cells from

subjects with asthma. In both NSA and SA sputum, there was a significant inverse relationship between LXA₄ and endogenous




Figure 5. Soluble epoxide hydrolase inhibition increases 15-epi-LXA₄ generation in severe asthma. Samples of (A) whole blood and (B) bronchoalveolar lavage fluids (BALF) were obtained from subjects with severe asthma and exposed to a soluble epoxide hydrolase inhibitor (sEHI) or vehicle before cell activation with A23187 (see METHODS). Materials from these incubations were extracted and 15-epi-LXA₄ was determined by ELISA. The change in 15-epi-LXA₄ is expressed as an increase relative to control samples exposed only to vehicle (whole blood, 74.0 ± 40.6 pg 15-epi-LXA₄/ml; BALF, 77.8 ± 55.4 pg 15-epi-LXA₄/ml). Results are expressed as mean \pm SEM ($n \geq 6$). * $P < 0.05$ in comparison with samples activated with A23187 in the absence of sEHI by one-way analysis of variance. (C) 15-epi-LXA₄, 14,15-EET, and 14,15-DHET were determined by ELISA. The ratio of 14,15-DHET to 14,15-EET was calculated as an index of SEH activity. Values for each individual sample are shown for whole blood. Analysis was based on the Kruskal-Wallis analysis of variance followed by the Mann-Whitney test with Bonferroni correction (A and B). Spearman rank correlation test was used to compare the ranks of two continuous variables (C, $r = -0.39$, $P = 0.01$). APAU = 1-(1-acetylpiperidin-4-yl)-3-adamantanylurea; AUDA = 12-(3-adamantan-1-yl-ureido) dodecanoic acid; 14,15-DHET = 14,15-dihydroxy-eicosatrienoic acid; 14,15-EET = 14,15-epoxy-eicosatrienoic acid; LX = lipoxin; TPPU = 1-trifluoromethoxyphenyl-3-(1-propionylpiperidin-4-yl) urea; *t*-TUCB = *trans*-4-[4-[3-(4-trifluoromethoxy-phenyl)-ureido]-cyclohexyloxy]-benzoic acid.

sEH activity (14,15-DHET/14,15-EET ratio). Moreover, sEHs blocked conversion of 14,15-EET to 14,15-DHET and significantly increased 15-epi-LXA₄ levels by SA whole blood and BALF cells. For these experiments, four structurally

different sEHs were used. These sEHs are tight binding inhibitors that approach stoichiometric interaction with sEH. They are competitive and reversible inhibitors that have high target occupancy and a slow off rate (37). The consistent findings with

all four compounds indicate a class effect for sEH inhibition, rather than an off-target action of a single compound. Detailed information on the synthesis, physical properties, and pharmacokinetics of each of these sEHs has been published (22, 35–37). Together, these findings support sEH inhibition as a new pharmacologic mechanism for enhancing endogenous LX generation in SA.

To assess the potential therapeutic impact of sEHs and LXs in asthma, two disease-related functional responses were selected for preclinical studies of human cells and lung tissues: leukocyte recruitment for airway inflammation and cytokine-mediated bronchial reactivity. For leukocyte recruitment, LPAs were quantitated in whole blood. LPAs play important roles in secondary capture for leukocyte entry into inflamed tissues, including lung (26, 27, 38). Asthmatic responses are associated with platelet activation (39, 40) and increased circulating LPAs have been detected in asthma attacks and after allergen challenge (39, 41). These LPAs can also serve as important sources for bioactive lipid mediators, including LTs and LXs (25, 33).

Here, we found that LPAs were increased in asthma and related to disease severity. PAF can activate both leukocytes and platelets (42) and *in vitro* initiated a marked increase in LPAs. LXA₄ gave concentration-dependent inhibition of PAF-initiated LPAs. LXA₄ and 15-epi-LXA₄ inhibitory responses were similar; however, neither 14,15-EET nor 8-IP led to significant inhibition. During incubations designed for LX generation from endogenous sources (as in [33]), sEHs partially inhibited PAF-induced LPAs, the amplitude of which was related to induction of 15-epi-LXA₄ generation. These findings indicated that LPAs are useful biomarkers of asthma activity and severity and that LXs can regulate platelet activation and interaction with peripheral blood leukocytes. If sEHs increase vascular LX generation, then subsequent lung inflammation could be decreased by disrupting LPAs. Thus, LPAs might also serve as a useful indicator of sEH efficacy. In addition, these results also suggest that some of the antiinflammatory actions attributed to EETs may relate to LX generation and action. The potential value of LPAs as cellular biomarkers of asthma activity and severity requires further




Figure 6. Leukocyte-platelet aggregates are associated with asthma severity and inhibited by lipoxins (LXs) and soluble epoxide hydrolase inhibition. Leukocyte-platelet aggregates in peripheral venous blood were detected and enumerated by flow cytometry before and after incubation with either platelet activating factor (PAF, 5 μ M) or vehicle control (0.1% ethanol). (A) Representative histograms of leukocyte-platelet aggregates (as determined by cells with forward and side scatter properties of granulocytes staining with the platelet-specific molecule CD41) in blood from a subject with severe asthma (SA; top) and a healthy subject (bottom). Inset: Percentages of leukocyte-platelet aggregates are shown. (B) Leukocyte-platelet aggregates (%) in blood from healthy subjects (HC, open) or subjects with nonsevere asthma (NSA, striped) or SA (cross-striped) that were exposed to vehicle (white) or PAF (gray) ($n \geq 3$). (C) Samples of whole blood were exposed to LX₄ (1–100 nM, 5 min) before activation with PAF (5 μ M) in samples from HC, subjects with NSA, or SA ($n > 3$). (D) Samples of whole blood from HC were exposed (5 min) to either vehicle (0.1% ethanol) or 100 nM of LX₄, 15-epi-LX₄, 14,15-epoxy-eicosatrienoic acid (14,15-EET), or 8-isoprostane before activation with PAF (5 μ M), and leukocyte-platelet aggregates (%) were determined. (E and F) Samples of whole blood from HC were exposed to granulocyte-macrophage colony-stimulating factor (200 pM, 50 min) followed by select soluble epoxide hydrolase inhibitors (sEHI) and then activated (15 min) with fMLP (10^{-7} M) and thrombin (0.1 U/ml). Leukocyte-platelet aggregates (%) and levels of 15-epi-LX₄ were determined ($n \geq 3$) (see METHODS). (B–F) Data are expressed as mean \pm SEM. * $P < 0.05$ in comparison with HC, ** $P < 0.05$ in comparison with vehicle, *** $P < 0.05$ in comparison with PAF, # $P < 0.05$ in comparison with 1 nM. TPPU = 1-trifluoromethoxyphenyl-3-(1-propionylpiperidin-4-yl) urea; t-TUCB = *trans*-4-[4-[3-(4-trifluoromethoxy-phenyl)-ureido]-cyclohexyloxy]-benzoic acid.

investigation in a larger, independent clinical study.

To investigate a direct role for LXs on lung tissue, we chose to investigate its effects on increased bronchial contraction, which is a major feature of asthma and is a principal cause of symptomatic dyspnea (1). When

given to subjects with asthma by inhalation, LX₄ can block LTC₄-induced bronchoprovocation (15), and LX₄ and its stable analogs can dampen MCh-initiated bronchial hyperreactivity in mice (14). Because sEHI-mediated LX generation requires cell-cell interactions between

leukocytes and airway cells (23) and leukocytes were not present in the lung tissue sections, the direct impact of 15-epi-LX₄ on bronchial responses in human lung was determined via isometric tension measurements on human bronchi in an isolated organ bath system. Exposure




Figure 7. Effect of 15-epi-LXA $_4$ on tumor necrosis factor (TNF)- α -induced increases in bronchial contraction. (A) Representative traces of histamine (1 μ M) induced tension on human bronchi cultured for 48 hours in either control media (blue dashed line), media containing TNF- α (10 ng/ml, blue solid line), or media with TNF- α and 15-epi-LXA $_4$ (100 nM, green dotted line) (see METHODS). (B) Representative traces of U-46619 induced tension on human bronchi under the same experimental conditions as described in A. (C) The impact of 15-epi-LXA $_4$ (100 nM) on tension induced by methacholine (MCh) (1 μ M), histamine (1 μ M), or U-46619 (30 nM) was measured with TNF- α -exposed human bronchi from multiple donors. (D) Percent inhibition of the TNF- α -triggered bronchial responses was determined for 15-epi-LXA $_4$ and 14,15-epoxy-eicosatrienoic acid (14,15-EET) (100 nM). Values represent the mean \pm SD for $n = 13$. * $P < 0.05$ by analysis of variance. (E) Representative recording of the MCh response and the concentration-dependent relaxing effects induced by LXA $_4$ in control condition and (F) in a paired recording in the presence of the ALX/FPR2 receptor antagonist WRW4 (1 μ M). (G) Results are expressed as the mean and SD for the effect of cumulative addition of 15-epi-LXA $_4$ in the absence (control) or presence of WRW4 on the tension induced by MCh (1 μ M) on human bronchi. $n = 12$; * $P < 0.05$. LX = lipoxin.

to 15-epi-LXA₄ significantly reduced TNF- α -induced increases in bronchial contraction to a wide range of agonists, including MCh, histamine, and a thromboxane receptor agonist (U-46619). LXs principally mediate their actions as agonists at ALX/FPR2 and/or antagonists at CysLT1 receptors (29). Here, an ALX/FPR2-selective antagonist partially blocked the actions of 15-epi-LXA₄ and 14,15-EET, suggesting roles for both LX receptors in regulating bronchial responses. Together, these findings suggest that a sEH that increased 15-epi-LXA₄ in SA would have beneficial actions on bronchial relaxation in inflamed airways.

In summary, LX generation in the asthmatic airway is adversely impacted by oxidative stress and sEH activity, in particular in SA. Specific inhibitors of sEH

increased 14,15-EET and 15-epi-LXA₄ generation by SA whole blood and BALF cells, and 15-epi-LXA₄ decreased PAF-initiated LPAs and TNF- α -mediated increases in bronchial contraction, suggesting that sEHs or LX stable analogs could serve as a novel pro-resolving therapeutic strategy in SA. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Acknowledgment: *Members of the National Heart, Lung, and Blood Institute's Asthma Clinical Research Network:* Bill T. Ameredes (University of Texas Medical Branch, Galveston, TX), Eugene Bleeker (Wake Forest School of Medicine, Winston-Salem, NC), William J. Calhoun (University of Texas Medical Branch, Galveston, TX), Reuben Cherniack (National Jewish Health, Denver, CO), Vernon M. Chinchilli (Penn State Hershey College of Medicine, Hershey, PA), Timothy J. Craig (Penn State

Hershey College of Medicine, Hershey, PA), Loren Denlinger (University of Wisconsin, Madison, WI), Emily DiMango (New York-Presbyterian/Columbia University Medical Center, New York, NY), John Fahy (University of California San Francisco, San Francisco, CA), Nikolina Icitovic (Penn State Hershey College of Medicine, Hershey, PA), Tonya S. King (Penn State Hershey College of Medicine, Hershey, PA), Monica Kraft (Duke University School of Medicine, Durham, NC), Stephen C. Lazarus (University of California San Francisco, San Francisco, CA), Robert F. Lemanske (University of Wisconsin, Madison, WI), Stephen Peters (Wake Forest School of Medicine, Winston-Salem, NC), Joe Ramsdell (University of California San Diego, San Diego, CA), Robert A. Smith (National Heart, Lung, and Blood Institute, Bethesda, MD), Christine A. Sorkness (University of Wisconsin, Madison, WI), Stanley J. Szefer (National Jewish Health, Denver, CO), Michael J. Walter (Washington University School of Medicine in St. Louis, St. Louis, MO), and Stephen I. Wasserman (University of California San Diego, San Diego, CA)

References

- Fanta CH. Asthma. *N Engl J Med* 2009;360:1002–1014.
- Jarjour NN, Erzurum SC, Bleeker ER, Calhoun WJ, Castro M, Comhair SA, Chung KF, Curran-Everett D, Dweik RA, Fain SB, et al.; NHLBI Severe Asthma Research Program (SARP). Severe asthma: lessons learned from the National Heart, Lung, and Blood Institute Severe Asthma Research Program. *Am J Respir Crit Care Med* 2012;185:356–362.
- Barnes PJ. Corticosteroid resistance in airway disease. *Proc Am Thorac Soc* 2004;1:264–268.
- Moore WC, Meyers DA, Wenzel SE, Teague WG, Li H, Li X, D'Agostino R Jr, Castro M, Curran-Everett D, Fitzpatrick AM, et al.; National Heart, Lung, and Blood Institute's Severe Asthma Research Program. Identification of asthma phenotypes using cluster analysis in the Severe Asthma Research Program. *Am J Respir Crit Care Med* 2010;181:315–323.
- Levy BD, Vachier I, Serhan CN. Resolution of inflammation in asthma. *Clin Chest Med* 2012;33:559–570.
- Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med* 2012;18:716–725.
- Serhan CN. Resolution phase of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu Rev Immunol* 2007;25:101–137.
- Haworth O, Cernadas M, Yang R, Serhan CN, Levy BD. Resolvin E1 regulates interleukin 23, interferon-gamma and lipoxin A4 to promote the resolution of allergic airway inflammation. *Nat Immunol* 2008;9:873–879.
- Samuelsson B, Dahlén SE, Lindgren JA, Rouzer CA, Serhan CN. Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. *Science* 1987;237:1171–1176.
- Lee TH, Crea AE, Gant V, Spur BW, Marron BE, Nicolaou KC, Reardon E, Brezinski M, Serhan CN. Identification of lipoxin A4 and its relationship to the sulfidopeptide leukotrienes C4, D4, and E4 in the bronchoalveolar lavage fluids obtained from patients with selected pulmonary diseases. *Am Rev Respir Dis* 1990;141:1453–1458.
- Kazani S, Planagumà A, Ono E, Bonini M, Zahid M, Marigowda G, Wechsler ME, Levy BD, Israel E. Exhaled breath condensate eicosanoid levels associate with asthma and its severity. *J Allergy Clin Immunol* 2013;132:547–553.
- Levy BD, Bonnans C, Silverman ES, Palmer LJ, Marigowda G, Israel E; Severe Asthma Research Program, National Heart, Lung, and Blood Institute. Diminished lipoxin biosynthesis in severe asthma. *Am J Respir Crit Care Med* 2005;172:824–830.
- Vachier I, Bonnans C, Chavis C, Farce M, Godard P, Bousquet J, Chanez P. Severe asthma is associated with a loss of LX4, an endogenous anti-inflammatory compound. *J Allergy Clin Immunol* 2005;115:55–60.
- Levy BD, De Sanctis GT, Devchand PR, Kim E, Ackerman K, Schmidt BA, Szczeklik W, Drazen JM, Serhan CN. Multi-pronged inhibition of airway hyper-responsiveness and inflammation by lipoxin A(4). *Nat Med* 2002;8:1018–1023.
- Christie PE, Spur BW, Lee TH. The effects of lipoxin A4 on airway responses in asthmatic subjects. *Am Rev Respir Dis* 1992;145:1281–1284.
- Wu SH, Chen XQ, Liu B, Wu HJ, Dong L. Efficacy and safety of 15(R/S)-methyl-lipoxin A(4) in topical treatment of infantile eczema. *Br J Dermatol* 2013;168:172–178.
- Planagumà A, Kazani S, Marigowda G, Haworth O, Mariani TJ, Israel E, Bleeker ER, Curran-Everett D, Erzurum SC, Calhoun WJ, et al. Airway lipoxin A4 generation and lipoxin A4 receptor expression are decreased in severe asthma. *Am J Respir Crit Care Med* 2008;178:574–582.
- Smith KR, Pinkerton KE, Watanabe T, Pedersen TL, Ma SJ, Hammock BD. Attenuation of tobacco smoke-induced lung inflammation by treatment with a soluble epoxide hydrolase inhibitor. *Proc Natl Acad Sci USA* 2005;102:2186–2191.
- Eickmeier O, Seki H, Haworth O, Hilberath JN, Gao F, Uddin M, Croze RH, Carlo T, Pfeffer MA, Levy BD. Aspirin-triggered resolvin D1 reduces mucosal inflammation and promotes resolution in a murine model of acute lung injury. *Mucosal Immunol* 2013;6:256–266.
- Morin C, Sirois M, Echave V, Gomes MM, Rousseau E. Epoxyeicosatrienoic acid relaxing effects involve Ca²⁺-activated K⁺ channel activation and CPI-17 dephosphorylation in human bronchi. *Am J Respir Cell Mol Biol* 2007;36:633–641.
- Zhang D, Xie X, Chen Y, Hammock BD, Kong W, Zhu Y. Homocysteine upregulates soluble epoxide hydrolase in vascular endothelium in vitro and in vivo. *Circ Res* 2012;110:808–817.
- Morin C, Sirois M, Echavé V, Albadine R, Rousseau E. 17,18-Epoxyeicosatetraenoic acid targets PPAR γ and p38 mitogen-activated protein kinase to mediate its anti-inflammatory effects in the lung: role of soluble epoxide hydrolase. *Am J Respir Cell Mol Biol* 2010;43:564–575.
- Planagumà A, Pfeffer MA, Rubin G, Croze R, Uddin M, Serhan CN, Levy BD. Lovastatin decreases acute mucosal inflammation via 15-epi-lipoxin A4. *Mucosal Immunol* 2010;3:270–279.
- Morisseau C, Hammock BD. Measurement of soluble epoxide hydrolase (sEH) activity. *Current Protoc Toxicol* 2007;33:4.23.1–4.23.18.

25. Laidlaw TM, Kidder MS, Bhattacharyya N, Xing W, Shen S, Milne GL, Castells MC, Chhay H, Boyce JA. Cysteinyl leukotriene overproduction in aspirin-exacerbated respiratory disease is driven by platelet-adherent leukocytes. *Blood* 2012;119:3790–3798.
26. Looney MR, Nguyen JX, Hu Y, Van Ziffle JA, Lowell CA, Matthey MA. Platelet depletion and aspirin treatment protect mice in a two-event model of transfusion-related acute lung injury. *J Clin Invest* 2009; 119:3450–3461.
27. Zarbock A, Singbartl K, Ley K. Complete reversal of acid-induced acute lung injury by blocking of platelet-neutrophil aggregation. *J Clin Invest* 2006;116:3211–3219.
28. Morin C, Sirois M, Echave V, Gomes MM, Rousseau E. EET displays anti-inflammatory effects in TNF-alpha stimulated human bronchi: putative role of CPI-17. *Am J Respir Cell Mol Biol* 2008;38:192–201.
29. Gronert K, Martinsson-Niskanen T, Ravasi S, Chiang N, Serhan CN. Selectivity of recombinant human leukotriene D(4), leukotriene B(4), and lipoxin A(4) receptors with aspirin-triggered 15-epi-LXA(4) and regulation of vascular and inflammatory responses. *Am J Pathol* 2001;158:3–9.
30. Dworski R, Murray JJ, Roberts LJ II, Oates JA, Morrow JD, Fisher L, Sheller JR. Allergen-induced synthesis of F(2)-isoprostanes in atopic asthmatics. Evidence for oxidant stress. *Am J Respir Crit Care Med* 1999;160:1947–1951.
31. Horváth I, Hunt J, Barnes PJ, Alving K, Antczak A, Baraldi E, Becher G, van Beurden WJ, Corradi M, Dekhuijzen R, et al.; ATS/ERS Task Force on Exhaled Breath Condensate. Exhaled breath condensate: methodological recommendations and unresolved questions. *Eur Respir J* 2005;26:523–548.
32. Wood LG, Garg ML, Simpson JL, Mori TA, Croft KD, Wark PA, Gibson PG. Induced sputum 8-isoprostane concentrations in inflammatory airway diseases. *Am J Respir Crit Care Med* 2005;171:426–430.
33. Fiore S, Serhan CN. Formation of lipoxins and leukotrienes during receptor-mediated interactions of human platelets and recombinant human granulocyte/macrophage colony-stimulating factor-primed neutrophils. *J Exp Med* 1990;172:1451–1457.
34. Deng Y, Theken KN, Lee CR. Cytochrome P450 epoxygenases, soluble epoxide hydrolase, and the regulation of cardiovascular inflammation. *J Mol Cell Cardiol* 2010;48:331–341.
35. Morisseau C, Goodrow MH, Newman JW, Wheelock CE, Dowdy DL, Hammock BD. Structural refinement of inhibitors of urea-based soluble epoxide hydrolases. *Biochem Pharmacol* 2002;63:1599–1608.
36. Rose TE, Morisseau C, Liu JY, Inceoglu B, Jones PD, Sanborn JR, Hammock BD. 1-Aryl-3-(1-acylpiperidin-4-yl)urea inhibitors of human and murine soluble epoxide hydrolase: structure-activity relationships, pharmacokinetics, and reduction of inflammatory pain. *J Med Chem* 2010;53:7067–7075.
37. Lee KS, Morisseau C, Yang J, Wang P, Hwang SH, Hammock BD. Förster resonance energy transfer competitive displacement assay for human soluble epoxide hydrolase. *Anal Biochem* 2013;434:259–268.
38. Pitchford SC, Yano H, Lever R, Riffo-Vasquez Y, Ciferri S, Rose MJ, Giannini S, Momi S, Spina D, O'connor B, et al. Platelets are essential for leukocyte recruitment in allergic inflammation. *J Allergy Clin Immunol* 2003;112:109–118.
39. Gresele P, Dottorini M, Selli ML, Iannacci L, Canino S, Todisco T, Romano S, Crook P, Page CP, Nenci GG. Altered platelet function associated with the bronchial hyperresponsiveness accompanying nocturnal asthma. *J Allergy Clin Immunol* 1993;91:894–902.
40. Kornerup KN, Page CP. The role of platelets in the pathophysiology of asthma. *Platelets* 2007;18:319–328.
41. Kowal K, Pampuch A, Kowal-Bielecka O, DuBuske LM, Bodzenta-Łukaszyk A. Platelet activation in allergic asthma patients during allergen challenge with *Dermatophagoides pteronyssinus*. *Clin Exp Allergy* 2006;36:426–432.
42. Yost CC, Weyrich AS, Zimmerman GA. The platelet activating factor (PAF) signaling cascade in systemic inflammatory responses. *Biochimie* 2010;92:692–697.